

# STRONGLY CHROMATIC METRO DOMINATION OF $P_n$ , $C_n$ AND $P_n^2$

María del Pilar Gómez, Diego A. Muñoz

Department of Physics, Universidad de Chile, Santiago, Chile

## ABSTRACT

A dominating set  $D$  of a graph  $G(V, E)$  is called metro dominating set  $G$  if for every pair of vertices  $u, v$ , there exists a vertex  $w$  in  $D$  such that  $d(u, w) \neq d(v, w)$ . A metro dominating set  $D$  is called strongly chromatic metro dominating set if for every vertex  $v \in D$  is from the same color class. The minimum cardinality strongly chromatic metro dominating set is called strongly chromatic metro domination number and is denoted by  $SC\gamma_\beta$ . In this paper we find strongly chromatic metro domination number of path, cycles and square of a path.

**Keywords:** metric dimension, metro domination, strongly chromatic metro domination, power graph.

**AMS Mathematics Subject Classification (2010):**05C56.

## I. INTRODUCTION

Let  $G(V, E)$  be a simple, non-trivial, undirected and non-null graphs. A graph  $G$  is  $k$ -colorable if there exists a  $k$ -coloring of  $G$ . One of the fastest growing areas within graph theory is the study of domination and related problem. A subset  $D$  of  $V$  is said to be a dominating set of  $G$  if every vertex in  $V-D$  is adjacent to a vertex in  $D$ .

The minimum cardinality of a dominating set is called the domination number of  $G$  and is denoted by  $\gamma(G)$ . A subset  $D$  of  $V$  is said to be a dom-chromatic set if  $D$  is a dominating set and  $\chi(\langle D \rangle) = \chi(G)$ . The dom-chromatic number  $\gamma_{ch}(G)$  of  $G$  is the minimum cardinality of a dom-chromatic set.

In 1976 F. Harary and R.A. Melter [1] introduced the notation of metric dimension. A vertex  $x \in V(G)$  resolves a pair of vertices  $u, w \in V(G)$  if  $d(v, x) \neq d(w, x)$ . A set of vertices  $S \subseteq V(G)$  resolves  $G$  and  $S$  is a resolving set of  $G$ , if every pair of distinct vertices of  $G$  are resolved by same vertex in  $S$ . A resolving set  $S$  of  $G$  with minimum cardinality is a metric dimension of  $G$  denoted by  $\beta(G)$ .

A dominating set  $D$  of  $V(G)$  having a property that for each pair of vertices  $u, v$  there exist a vertex  $w$  in  $D$  such that  $d(u, w) \neq d(v, w)$  is called metro dominating set of  $G$  or simply MD-set. The minimum cardinality of a metro dominating set of  $G$  is called metro domination number of  $G$  and is denoted by  $\gamma_\beta(G)$ .

## II. DEFINITIONS

### 2.1 Metric dimension:

The metric dimension of a graph  $G$  is the minimum cardinality of a subset  $S$  of vertices such that all other vertices are uniquely determined by their distances to the vertices in  $S$ . It is denoted by  $\beta(G)$ .

### 2.2 Domination:

Let  $G(V, E)$  be a graph. A subset of vertices  $D \subseteq V$  is called a dominating set ( $\gamma$ -set) if every vertex in  $V-D$  adjacent to atleast one vertex in  $D$ . The minimum cardinality of a dominating set is called the domination number of the graph  $G$  and is denoted by  $\gamma(G)$ .

### 2.3 Locating domination:

A dominating set  $D$  is called a locating dominating set or simply LD-set if for each pair of vertices  $u, v \in V-D$ ,  $ND(u) \neq ND(v)$  where  $ND(u) = N(u) \cap D$ . The minimum cardinality of an LD-set of the graph  $G$  is called the locating domination number of  $G$  denoted by  $\gamma_L(G)$ .

### 2.4 Metro domination:

A dominating set  $D$  of  $V(G)$  having the property that for each pair of vertices  $u, v$  there exists a vertex  $w$  in  $D$  such that  $d(u, w) \neq d(v, w)$  is called metro dominating set of  $G$  or simply MD-set. The minimum cardinality of a metro dominating set of  $G$  is called metro domination number of  $G$  and is denoted by  $\gamma_\beta(G)$ .

**2.5 Chromatic number:**

The minimum number of colors required for a proper coloring of  $G$  is called chromatic number of  $G$  and is denoted by  $\chi(G)$ .

**2.6 Chromatic domination:**

A subset  $D$  of  $V$  is said to be a dom-chromatic set if  $D$  is a dominating set and  $\chi(\langle D \rangle) = \chi(G)$ . The dom-chromatic number  $\gamma_{ch}(G)$  of  $G$  is the minimum cardinality of a dom-chromatic set.

**III. SOME KNOWN RESULTS**

In this section we mention some of the known result on metric dimension, domination, metro domination.

**Theorem 3.1.** (Harary and Melter [1]) The metric dimension of a non trivial complete graph of order  $n$  is  $n-1$ .

**Theorem 3.2.** (Khuller, Raghavachari, Rosenfeld [4]) The metric dimension of a graph  $G$  is 1 if and only if  $G$  is a path.

**Theorem 3.3.** (Harary and Melter [1]) The metric dimension of a complete bipartite graph  $K_{m,n}$  is  $m+n-2$ .

**Theorem 3.4.**[5] The metro domination number of a graph  $G$  is  $\left\lceil \frac{n}{5} \right\rceil$  if and only if  $G$  is a cycle.

**Theorem 3.5.**[5] Let  $G$  be a graph on  $n$  vertices. Then  $\gamma_\beta(G) = n-1$  if and only if  $G$  is  $K_n$  or  $K_{1,n-1}$  for  $n \geq 1$ .

**Theorem 3.6.** [5] For any integer  $n$ ,  $\gamma_\beta(P_n) = \left\lceil \frac{n}{3} \right\rceil$ .

**Remark 3.7.** For any connected graph  $G$ ,  $\gamma_\beta(G) \geq \max\{\gamma(G), \beta(G)\}$ .

**Remark 3.8.** For any integer  $n > 3$ ,  $\chi(C_n) = \begin{cases} 3 & \text{for } n \text{ odd} \\ 2 & \text{for } n \text{ even} \end{cases}$

**Remark 3.9.** For any integer  $n > 1$ ,  $\chi(P_n) = 2$ .

**Lemma 3.10.** [9] Let  $G = P_n^2$ ,  $n > 3$ . Then  $\dim(G) = 2$ .

**Theorem 3.11.**[7] For every  $n \geq 1$ ,  $\gamma_\beta(P_n^2) = \left\lceil \frac{n}{5} \right\rceil$ .

**Theorem 3.12.** [2] For any integer  $n \geq 3$ ,  $\gamma_\beta(P_n^2) = \begin{cases} 2 & \text{if } 3 \leq n \leq 7 \\ 3 & \text{if } 8 \leq n \leq 10 \\ \left\lceil \frac{n}{5} \right\rceil & \text{if } n \geq 11 \end{cases}$

**Remark 3.13.** For any integer  $n \geq 3$ ,  $\chi(P_n^2) = 3$ .

**IV. MAIN RESULTS**

**Theorem 4.1.** For any integer  $n \geq 4$ ,  $SC\gamma_\beta(P_n) = \left\lceil \frac{n-1}{2} \right\rceil$ .

**Proof:** By theorem 3.2  $\beta(P_n) = 1$  and by remark 3.9  $\chi(P_n) = 2$ , clearly we have  $\left\lceil \frac{n}{2} \right\rceil$  vertices of one color class and remaining  $\left\lfloor \frac{n}{2} \right\rfloor$  vertices of other color class. Hence we have choice of either  $\left\lceil \frac{n}{2} \right\rceil$  or  $\left\lfloor \frac{n}{2} \right\rfloor$  vertices for dominating set  $D$

whose vertices are from the same color class. For even  $n$ ,  $\frac{n}{2}$  vertices of same color class dominates the remaining  $\frac{n}{2}$  vertices. For odd  $n$ ,  $\frac{n-1}{2}$  vertices of same color class will dominates the remaining vertices and hence  $SC\gamma_\beta(P_n) \geq \left\lfloor \frac{n-1}{2} \right\rfloor$

(1)

To prove the reverse inequality, we define a strongly chromatic dominating set  $D = \{v_{2i} / 1 \leq i \leq \left\lfloor \frac{n}{2} \right\rfloor\}$  of cardinality  $\left\lfloor \frac{n-1}{2} \right\rfloor$ . We note that  $D$  acts as a dominating set also as a resolving set and each  $v_i \in D$  is from the same color class and hence  $SC\gamma_\beta(P_n) \leq \left\lfloor \frac{n-1}{2} \right\rfloor$

(2)

from (1) and (2)

$$SC\gamma_\beta(P_n) = \left\lfloor \frac{n-1}{2} \right\rfloor.$$

**Theorem 4.2.** For any integer  $n \geq 5$ ,  $SC\gamma_\beta(C_n) = \left\lfloor \frac{n-1}{2} \right\rfloor$ .

**Proof:** By the result  $\beta(C_n) = 2$  and by remark 3.8,  $\chi(C_n) = \begin{cases} 3 & \text{for } n \text{ odd} \\ 2 & \text{for } n \text{ even} \end{cases}$ , clearly we have  $\frac{n}{2}$  vertices of one color class and remaining  $\frac{n}{2}$  vertices of other color class for even  $n$  and  $\left\lfloor \frac{n}{2} \right\rfloor$  vertices of one color class and other  $\left\lfloor \frac{n}{2} \right\rfloor$  vertices of second color class and remaining one vertex of third color class for odd  $n$ . Hence we have choice of  $\frac{n}{2}$  vertices for dominating set  $D$  such that each  $v_i \in D$  are from the same color class. For even cycle,  $\frac{n}{2}$  vertices of same color class dominate the remaining  $\frac{n}{2}$  vertices. For odd cycle,  $\frac{n-1}{2}$  vertices of same color class will dominate the remaining vertices and hence  $SC\gamma_\beta(C_n) \geq \left\lfloor \frac{n-1}{2} \right\rfloor$

(1)

To prove the reverse inequality, we define a strongly chromatic dominating set  $D = \{v_{2i-1} / 1 \leq i \leq \left\lfloor \frac{n}{2} \right\rfloor\}$  of cardinality  $\left\lfloor \frac{n-1}{2} \right\rfloor$ , which also acts as a resolving set and each  $v_i \in D$  is from the same color class and hence  $SC\gamma_\beta(C_n) \leq \left\lfloor \frac{n-1}{2} \right\rfloor$

(2)

from (1) and (2)

$$SC\gamma_\beta(C_n) = \left\lfloor \frac{n-1}{2} \right\rfloor.$$

**Theorem 4.3.** For any integer  $n \leq 9$ ,  $SC\gamma_\beta(P_n^2) = \left\lfloor \frac{n}{3} \right\rfloor$ .

**Proof:** By lemma 3.10,  $\dim(P_n^2) = 2$  for  $n > 3$ . Also by Theorem 3.11  $\gamma_\beta(P_n^2) = \left\lfloor \frac{n}{5} \right\rfloor$ ,  $n \geq 11$  here each  $\left\lfloor \frac{n}{5} \right\rfloor$  vertices of metro dominating set are not from the same color class. By remark 3.13,  $\chi(P_n^2) = 3$ ,  $n \geq 3$  if we label  $v_1$  of  $P_n^2$  by color 1 and  $v_2$  by color 2 and  $v_3$  by color 3 and continuing the coloring, we get  $\left\lfloor \frac{n}{3} \right\rfloor$  vertices of color class 1,  $\left\lfloor \frac{n+1}{3} \right\rfloor$  vertices of color class 2 and  $\left\lfloor \frac{n}{3} \right\rfloor$  vertices of color class 3. Hence we have a choice of  $\left\lfloor \frac{n}{3} \right\rfloor$  or  $\left\lfloor \frac{n+1}{3} \right\rfloor$  or  $\left\lfloor \frac{n}{3} \right\rfloor$  vertices for strongly chromatic metro dominating set minimum among these  $\left\lfloor \frac{n}{3} \right\rfloor$  is minimum and hence

$$SC\gamma_\beta(P_n^2) \geq \left\lfloor \frac{n}{3} \right\rfloor$$


(1)

To prove the reverse inequality, we defined the strongly chromatic metro dominating set as  $D = \{v_{3i} / 1 \leq i \leq \left\lfloor \frac{n}{3} \right\rfloor\}$  of cardinality  $\left\lfloor \frac{n}{3} \right\rfloor$ . We note that  $D$  is a dominating set also acts as a resolving set and each  $v_i \in D$  are all from the same color class and hence  $SC\gamma_\beta(P_n^2) \leq \left\lfloor \frac{n}{3} \right\rfloor$  (2)

from (1) and (2)

$$SC\gamma_\beta(P_n^2) = \left\lceil \frac{n}{3} \right\rceil.$$

**EXAMPLE:**



$$D_1 = \{v_3\}$$

$D_1$  is a dominating set but not resolving set.

$$D_2 = \{v_1, v_5\}$$

$D_2$  is a dominating set also resolving set but both vertices are not from same color class. Hence it is not a strongly chromatic metro domination.

Hence  $P_5^2$  is not a strongly chromatic metro domination.

## REFERENCE

1. F. Harary and R.A. Melter, on the metric dimension of graphs, *Ars Combinatoria*, 2(1976) 191-195.
2. G.C. Basavaraju, M. Vishukumar and P. Raghunath, Metro domination of Square Path, *Annals of Pure and Applied Mathematics* vol.14, No. 3, 2017, 539-545.
3. S. Balamurugan, P. Aristotle, V. Swaminathan and G. Prabakaran, on graphs whose neighbourhood chromatic domination number is two, *International Journal of Pure and Applied Mathematics*, Volume 119, No. 15, 2018, 213-224.
4. S. Khuller, B. Raghavachari and A. Rosenfeld, Landmarks in graph, *Discrete Appl. Math.*, 70 (3) (1996) 217-229.
5. P. Raghunath and B. Sooryanarana, Metro domination number of a graph, *Twentieth annual conference of Ramanujan Mathematics Society*, July 25-30 (2005) University of Calicut.
6. S. Lakshminarayana and M. Vishukumar, on the  $k$ -Metro domination number of Paths, *Annals of Pure and Applied Mathematics*, Vol. 14, No. 3, 2017, 593-600.
7. M. Alishahi and S.H. Shalmaee, Domination number of Square of Cartesian product of Cycles, *Open Journal of Discrete Mathematics*, 5(2015) 88-94.
8. G.C. Basavaraju, M. Vishukumar and P. Raghunath, Metro domination of Square Cycle, *International Journal of Mathematics and its Applications*, Volume 5, Issue 4-E(2017), 641-645.
9. M. M. AlHoli, O.A. Abughneim and H.A. Ezech, Metric dimension of Path related graphs, *Global Journal of Pure and Applied Mathematics*, 13 (2) (2017) 149-157.